

Regence

Medical Policy Manual

Transplant, Policy No. 19

Uterus Transplant

Effective: September 1, 2023

Next Review: May 2024

Last Review: July 2023

IMPORTANT REMINDER

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

DESCRIPTION

Absolute uterine factor infertility is a condition in which an individual is unable to achieve pregnancy due to an absent or non-functioning uterus. Uterus transplantation is a complex, multi-stage process involving a living or deceased donor, recipient, and genetic partner.

MEDICAL POLICY CRITERIA

Uterus transplantation is considered **investigational** for all indications.

NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.

CROSS REFERENCES

None

BACKGROUND

ABSOLUTE UTERINE FACTOR INFERTILITY

Absolute uterine factor infertility (AUF) refers to infertility that is attributable to an absent or non-functional uterus due to congenital, surgical, anatomical, or acquired factors that prevent embryo implantation and term pregnancy. AUF is estimated to impact 1 in 500 females of childbearing age.^[1, 2]

Uterine agenesis or Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome results in the congenital absence of the uterus or presence of a rudimentary solid bipartite uterus. MRKH syndrome accounts for less than 3% of all müllerian malformations with an estimated prevalence of 1 in 4,500 females.^[3, 4] Individuals with MRKH syndrome type I present with two kidneys and are considered ideal candidates for uterine transplantation. Individuals with MRKH syndrome type II presenting with a single kidney have a higher risk of medication-induced nephrotoxicity and associated obstetric complications (e.g., severe preeclampsia).^[5]

Hysterectomy is the most common cause of acquired AUF, with 240,000 procedures taking place in females under age 44 in the United States.^[6] In one clinical trial screening study of 239 individuals at the Cleveland Clinic, indications for uterus transplantation included prior hysterectomy (64%) and congenital anomalies (32%). Among individuals with prior hysterectomy, 50% were performed for benign indications, 25% for malignancy, and 25% for obstetric complications.^[7]

UTERUS TRANSPLANTATION

Uterus transplantation may provide a unique fertility restoration option for individuals desiring to carry and birth a child.^[8] Uterus transplantation is a complex, multi-stage process involving a living or deceased donor, recipient, and genetic partner. Once screening and consent is established for all involved parties, in-vitro fertilization is performed prior to transplantation to ensure fertilization and normal embryo development.^[9] The transplantation surgery involves radical hysterectomy in the donor to ensure long vascular pedicles for transplantation;^[10] however, several cases of robot-assisted laparoscopic approaches have been reported.^[11, 12] An advantage of uterus procurement in a deceased donor involves freedom to transect ureters, but this convenience is balanced by the potential for prolonged uterus ischemic time.^[13] The surgical approach in the recipient is dictated by underlying pelvic anatomy which may be impacted by AUF etiology. For example, in individuals with Asherman syndrome, a traditional total hysterectomy must first be performed in the recipient. Immunosuppression is initiated at the time of transplantation and protocol and for-cause cervical biopsies enable monitoring for organ rejection.^[14, 15] After 6 to 12 months of immunosuppression, embryo transfer, pregnancy, and cesarean delivery may follow. When childbearing has been deemed complete, the transplanted uterus is removed to avoid lifelong immunosuppression. Thus, uterus transplantation is the first form of organ transplantation intended to be temporary.^[1, 9]

The first human uterus transplant was performed in 2000 in Saudi Arabia with a 46-year-old living donor and 26-year-old recipient that had acquired AUF due to hysterectomy for prior post-partum hemorrhage. Due to the development of acute vascular thrombosis at three months post-transplant, graft hysterectomy was required.^[16] The first successful live birth occurred in 2014 in Sweden in a 35-year-old recipient with MRKH syndrome via a living, 61 year old, two-parous donor. The recipient was admitted with preeclampsia at 31 weeks, and a healthy male child was born five days later via cesarean delivery.^[17] The first live birth in the United States occurred in 2017 in a 29 year old recipient with MRKH syndrome via a living, 32-year-old, two-parous donor.^[18] According to the Organ Procurement and Transplantation

Network (OPTN), 35 uterus transplants have been performed in the United States via 13 deceased and 22 living donors as of March 2022.^[19]

Literature has explored the implications of uterus transplantation in transgender women, identifying several theoretical medical issues in genetic males meriting further investigation. These include creation of adequate de novo uterine vascularization, administration of appropriate hormone replacement therapy, and placement of the donor uterus in a nongynecoid uterus.^[20, 21]

REGULATORY STATUS

Solid organ transplants are a surgical procedure and, as such, are not subject to regulation by the U.S. Food and Drug Administration (FDA).

The FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation Title 21, parts 1270 and 1271. Solid organs used for transplantation are subject to these regulations.

Restorative or life-enhancing uterine vascularized composite allograft (VCA) procurement and transplantation falls under the oversight of the Organ Procurement and Transplantation Network (OPTN).^[22]

EVIDENCE SUMMARY

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome, defined as length of life, quality of life, and ability to function including benefits and harms. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. Randomized controlled trials (RCTs) are preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

There are no RCTs directly comparing uterus transplant with alternatives. Systematic reviews are based on case series. Studies comparing surgical technique, infection prophylaxis, and immunosuppressive regimens are not germane to this evidence review.

SYSTEMATIC REVIEWS

Brannstrom (2021) published a systematic review of all published clinical uterus transplantation data and major interim results from 2000 through 2019.^[1] Of 62 uterus transplants identified for the review, the overall technical success rate defined as subsequent regular menstruation, was 76%. Technical success rates for living and deceased donor procedures were 78% and 64%, respectively. Rates of serious postsurgical complications were 18% for living donors and 19% for recipients. Most uterus transplantation procedures to date have involved living donors (51/62, 82%). Complications in living donors have included ureteric laceration, urinary bladder hypotonia, unplanned bilateral oophorectomy, vaginal dehiscence, fecal impaction, and unilateral pyelonephritis and hydronephrosis. Postoperative complications

in recipients have included vaginal anastomotic stenosis and treatable episodes of minor to severe graft rejection.

The cumulative live birth rate per transplant attempt, and per surgically successful uterus transplant is estimated to be >60% and >80%, respectively, as based on 24 published live birth accounts from interim data. High rates of preterm birth (19/24, 80%) and respiratory distress syndrome in the newborn (9/24, 38%) have been observed across cases. Obstetric complications have included preeclampsia, gestational hypertension, and several cases of placenta previa and gestational diabetes. Newborns had an Apgar score of 8 or higher at five minutes. One minor malformation in a female newborn involving an anteriorly caudally displaced urethra was reported, which was surgically corrected at 11 months. The reviewers concluded that "the modest success rate and the fairly high complication rate among [living donors], indicate that further research and development under strict governance are needed before this option should be widely offered."

Case Series

Characteristics and interim results from select case series are summarized in Tables 1 and 2.

Table 1. Summary of Key Case Series Characteristics

Study	Country (Years)	LD Criteria	Recipient Criteria	Participants
Johannesson (2021); ^[23] Putman (2021) ^[24]	United States (2016-2019)	NR	Women with AUFI and intact native ovaries and of childbearing age 20 to 35, negative history of or prior vaccination for HPV, and meets physiological criteria	Median age, 30 years (range, 20 to 35); 17 MRKH type I 1 MRKH type II 2 prior hysterectomy for leiomyoma(s) Mean donor age, NR 18 LD UTx; 2 DD UTx
Fronek (2021) ^[25]	Czech Republic (2016-2018)	Female 18 to 60 years of age, ≤4 childbirths, ≤1 cesarean section, good general health	Female 18 to 40 years of age, AUFI based on congenital or acquired uterus absence, desire for a child, having a male partner, and good general health	Mean recipient age, 28±3 years; 9 MRKH type I; 1 MRKH type II; Mean donor age, 46±14 years; 5 LD UTx (all related); 5 DD UTx; 5 postmenopausal; 2 nulliparous

AUFI: absolute uterine factor infertility; DD: deceased donor; HPV: human papillomavirus; LD: living donor; MRKH: Mayer-Rokitansky-Küster-Hauser syndrome; NR: not reported; UTx: uterus transplant.

Table 2. Summary of Key Case Series Results

Study	Survival	Embryo Transfers, total (range)	Clinical Pregnancy, total (n)	Live Births, total (n)	Live Birth Success Rate	Complications
Johannesson (2021); ^[23] Putman (2021) ^[24]	Graft: 14/20 (70% at 3 years)	30 (1 to 6)	19 (14)	13 (12); 11 LD and 1 DD; 10 MRKH	Overall: 60%; With surgical	acute rejection, gestational hypertension, preeclampsia,

Study	Survival	Embryo Transfers, total (range)	Clinical Pregnancy, total (n)	Live Births, total (n)	Live Birth Success Rate	Complications
				type I; 1 MRKH type II; 1 prior hysterectomy	success: 86%	gestational diabetes mellitus, placenta previa, preterm delivery
Fronek (2021) ^[25]	Graft: 7/10 (70% at 1 year); Recipient: 10/10 (100% at 2 years)	40 (4 to 11)	7 (5)	3 (3); 2 LD and 1 nulliparous DD	Overall: 30%; With surgical success: 43%	vaginal stenosis, leukopenia, UTI, acute rejection, CMV replication, graft HSV infection, <i>C. difficile</i> infection; HLA mismatch, CKD

CKD: chronic kidney disease; CMV: cytomegalovirus; DD: deceased donor; HLA: human leukocyte antigen; HSV: herpes simplex virus; LD: living donor; MRKH: Mayer-Rokitansky-Küster-Hauser syndrome; UTI: urinary tract infection.

Section Summary

Case series of uterus transplantation for AUFI have predominantly enrolled individuals with MRKH syndrome type I. A systematic review of interim trial data has reported live birth success estimates exceeding 60% overall and 80% among transplant attempts with surgical success. Slightly higher technical success rates have been reported for living donor compared to deceased donor procedures (78% vs. 64%, respectively). Rates of serious complications are high among both recipients (19%) and living donors (18%). High rates of preterm birth (80%) and episodes of acute respiratory distress syndrome in the newborn have been reported. Long-term health outcomes in children born via uterus transplantation and recipients following graft hysterectomy continue to accumulate in ongoing trials.

PRACTICE GUIDELINE SUMMARY

AMERICAN COLLEGE OF OBSTETRICIANS AND GYNECOLOGISTS

In 2018 (reaffirmed 2020), the American College of Obstetricians and Gynecologists (ACOG) Committee on Adolescent Health Care issued a Committee Opinion (Number 728) on the diagnosis, management, and treatment of müllerian agenesis.^[26] Regarding future fertility options, the opinion states that while live births have resulted from uterine transplantation, "given limited data, this procedure currently is considered experimental and is not widely available."

AMERICAN SOCIETY FOR REPRODUCTIVE MEDICINE

In 2018, the American Society for Reproductive Medicine (ASRM) issued a position statement recognizing uterus transplantation as the first successful medical treatment for absolute uterine

factor infertility, emphasizing its experimental nature.^[27] The statement recommends that the procedure should be performed within an Institutional Review Board-approved research protocol, with recommendations for the composition of "well-coordinated and multidisciplinary" uterus transplantation teams and suggested recipient inclusion and exclusion criteria.

SUMMARY

There is not enough evidence to show that uterus transplantation can improve health outcomes for patients. Further study is necessary to increase success rates, decrease complications and preterm births, and assess long-term outcomes in recipients and their children. In addition, there are no evidence-based clinical practice guidelines that recommend this procedure. Therefore, uterus transplant is considered investigational for all indications.

REFERENCES

1. Brannstrom M, Belfort MA, Ayoubi JM. Uterus transplantation worldwide: clinical activities and outcomes. *Curr Opin Organ Transplant*. 2021;26(6):616-26. PMID: 34636769
2. Hellstrom M, El-Akouri RR, Sihlbom C, et al. Towards the development of a bioengineered uterus: comparison of different protocols for rat uterus decellularization. *Acta Biomater*. 2014;10(12):5034-42. PMID: 25169258
3. Grimbizis GF, Camus M, Tarlatzis BC, et al. Clinical implications of uterine malformations and hysteroscopic treatment results. *Hum Reprod Update*. 2001;7(2):161-74. PMID: 11284660
4. Folch M, Pigem I, Konje JC. Mullerian agenesis: etiology, diagnosis, and management. *Obstet Gynecol Surv*. 2000;55(10):644-9. PMID: 11023205
5. Garg AX, Nevis IF, McArthur E, et al. Gestational hypertension and preeclampsia in living kidney donors. *N Engl J Med*. 2015;372(2):124-33. PMID: 25397608
6. Brett KM, Higgins JA. Hysterectomy prevalence by Hispanic ethnicity: evidence from a national survey. *Am J Public Health*. 2003;93(2):307-12. PMID: 12554591
7. Arian SE, Flyckt RL, Farrell RM, et al. Characterizing women with interest in uterine transplant clinical trials in the United States: who seeks information on this experimental treatment? *Am J Obstet Gynecol*. 2017;216(2):190-91. PMID: 27865979
8. Jarvholm S, Enskog A, Hammarling C, et al. Uterus transplantation: joys and frustrations of becoming a 'complete' woman-a qualitative study regarding self-image in the 5-year period after transplantation. *Hum Reprod*. 2020;35(8):1855-63. PMID: 32619006
9. Malasevskaia I, Al-Awadhi AA. A New Approach for Treatment of Woman With Absolute Uterine Factor Infertility: A Traditional Review of Safety and Efficacy Outcomes in the First 65 Recipients of Uterus Transplantation. *Cureus*. 2021;13(1):e12772. PMID: 33614361
10. Johannesson L, Diaz-Garcia C, Leonhardt H, et al. Vascular pedicle lengths after hysterectomy: toward future human uterus transplantation. *Obstet Gynecol*. 2012;119(6):1219-25. PMID: 22617587
11. Wei L, Xue T, Tao KS, et al. Modified human uterus transplantation using ovarian veins for venous drainage: the first report of surgically successful robotic-assisted uterus

- procurement and follow-up for 12 months. *Fertil Steril*. 2017;108(2):346-56 e1. PMID: 28778283
12. Ayoubi JM, Carbonnel M, Pirtea P, et al. Laparotomy or minimal invasive surgery in uterus transplantation: a comparison. *Fertil Steril*. 2019;112(1):11-18. PMID: 31277761
 13. Gauthier T, Piver P, Pichon N, et al. Uterus retrieval process from brain dead donors. *Fertil Steril*. 2014;102(2):476-82. PMID: 24837613
 14. Molne J, Broecker V, Ekberg J, et al. Monitoring of Human Uterus Transplantation With Cervical Biopsies: A Provisional Scoring System for Rejection. *Am J Transplant*. 2017;17(6):1628-36. PMID: 27868389
 15. Balko J, Novackova M, Skapa P, et al. Histopathological examination of the ectocervical biopsy in non-transplanted uteri: A study contributing to the provisional scoring system of subclinical graft rejection after uterus transplantation. *Acta Obstet Gynecol Scand*. 2022;101(1):37-45. PMID: 34693986
 16. Fageeh W, Raffa H, Jabbad H, et al. Transplantation of the human uterus. *Int J Gynaecol Obstet*. 2002;76(3):245-51. PMID: 11880127
 17. Brannstrom M, Johannesson L, Bokstrom H, et al. Livebirth after uterus transplantation. *Lancet*. 2015;385(9968):607-16. PMID: 25301505
 18. Testa G, McKenna GJ, Gunby RT, Jr., et al. First live birth after uterus transplantation in the United States. *Am J Transplant*. 2018;18(5):1270-74. PMID: 29575738
 19. Organ Procurement and Transplantation Network (OPTN). National data: Transplants by Donor Type [GU: Uterus]. March 2022. [cited 07/05/2023]. 'Available from:' <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/#>.
 20. Lefkowitz A, Edwards M, Balayla J. Ethical considerations in the era of the uterine transplant: an update of the Montreal Criteria for the Ethical Feasibility of Uterine Transplantation. *Fertil Steril*. 2013;100(4):924-6. PMID: 23768985
 21. Jones BP, Rajamanoharan A, Vali S, et al. Perceptions and Motivations for Uterus Transplant in Transgender Women. *JAMA Netw Open*. 2021;4(1):e2034561. PMID: 33471119
 22. Organ Procurement and Transplantation Network (OPTN). Vascular composite allograft. n.d. [cited 07/05/2023]. 'Available from:' <https://optn.transplant.hrsa.gov/professionals/by-organ/vascular-composite-allograft>.
 23. Johannesson L, Testa G, Putman JM, et al. Twelve Live Births After Uterus Transplantation in the Dallas UtErus Transplant Study. *Obstet Gynecol*. 2021;137(2):241-49. PMID: 33416285
 24. Putman JM, Zhang L, Gregg AR, et al. Clinical pregnancy rates and experience with in vitro fertilization after uterus transplantation: Dallas Uterus Transplant Study. *Am J Obstet Gynecol*. 2021;225(2):155 e1-55 e11. PMID: 33716072
 25. Fronek J, Kristek J, Chlupac J, et al. Human Uterus Transplantation from Living and Deceased Donors: The Interim Results of the First 10 Cases of the Czech Trial. *J Clin Med*. 2021;10(4). PMID: 33557282
 26. Committee on Adolescent Health C. ACOG Committee Opinion No. 728: Mullerian Agenesis: Diagnosis, Management, And Treatment. *Obstet Gynecol*. 2018;131(1):e35-e42. PMID: 29266078
 27. Practice Committee of the American Society for Reproductive Medicine. Electronic address aao, Practice Committee of the American Society for Reproductive M. American Society for Reproductive Medicine position statement on uterus transplantation: a committee opinion. *Fertil Steril*. 2018;110(4):605-10. PMID: 30196945

CODES

Codes	Number	Description
CPT	0664T	Donor hysterectomy (including cold preservation); open, from cadaver donor
	0665T	Donor hysterectomy (including cold preservation); open, from living donor
	0666T	Donor hysterectomy (including cold preservation); laparoscopic or robotic, from living donor
	0667T	Recipient uterus allograft transplantation from cadaver or living donor
	0668T	Backbench standard preparation of cadaver or living donor uterine allograft prior to transplantation, including dissection and removal of surrounding soft tissues and preparation of uterine vein(s) and uterine artery(ies), as necessary
	0669T	Backbench reconstruction of cadaver or living donor uterus allograft prior to transplantation; venous anastomosis, each
	0670T	Backbench reconstruction of cadaver or living donor uterus allograft prior to transplantation; arterial anastomosis, each
HCPCS	None	

Date of Origin: May 2022